VOL. 603, NOS. 1+2 JOURNAL OF CHROMATOGRAPHY JUNE 19, 1992

CONTENTS

(Abstracts/Contents Lists published in Analytical Abstracts, Biochemical Abstracts, Biological Abstracts, Chemical Abstracts, Chemical Titles, Chromatography Abstracts, Current Contents/Life Sciences, Current Contents/Physical, Chemical & Earth Sciences, Deep-Sea Research/Part B: Oceanographic Literature Review, Excerpta Medica, Index Medicus, Mass Spectrometry Bulletin, PASCAL-CRNS, Referativnyi Zhurnal, Research Alert and Science Citation Index)

REGULAR PAPERS

Column Liquid Chromatography

Comparison of the various kinetic models of non-linear chromatography
by S. Golshan-Shirazi and G. Guiochon (Knoxville and Oak Ridge, TN, USA) (Received February 12th, 1992) . . . 1

Comparison between the hodograph transform method and frontal chromatography for the measurement of binary competitive adsorption isotherms
by Z. Ma and G. Guiochon (Knoxville and Oak Ridge, TN, USA) (Received February 5th, 1992) 13

Batch dynamic adsorption of dipeptides onto reversed-phase silica gel
by T. S. Nguyen, S.-G. Hu and D. D. Do (Brisbane, Australia) (Received March 10th, 1992) 27

The *S* index in the retention equation in reversed-phase high-performance liquid chromatography
by N. Chen, Y. Zhang and P. Lu (Dalian, China) (Received February 11th, 1992) 35

Dispersion in stacked-membrane chromatography
by D. D. Frey, R. Van de Water and B. Zhang (New Haven, CT, USA) (Received February 25th, 1992) 43

Reduction in large-scale adsorption chromatography unit costs by use of 30-Å Sorbsil C30 silica
by I. Chappell and P. Baines (Warrington, UK) and P. K. Carpenter (Bracknell, UK) (Received January 31st, 1992) . 49

Solvent–stationary phase interactions in normal bonded phase high-performance liquid chromatographic columns. I. Investigation of system peaks in amino bonded phase columns
by C. W. Hsu and W. T. Cooper (Tallahassee, FL, USA) (Received February 25th, 1992) 63

Support for liquid–liquid partition chromatography in aqueous two-phase systems: a comparison of Superdex and LiParGel
by C. Wingren, U.-B. Hansson and K. Andersson (Lund, Sweden) (Received February 17th, 1992) 73

Systematic study on the resolution of derivatized amino acids enantiomers on different cyclodextrin-bonded stationary phases
by S. H. Lee, A. Berthod and D. W. Armstrong (Rolla, MO, USA) (Received February 21st, 1992) 83

Protein purification using combined streptavidin (or avidin)-Sepharose and thiopropyl-Sepharose affinity chromatography
by F. Desarnaud, J. Marie, R. Larguier, C. Lombard, S. Jard and J.-C. Bonnafous (Montpellier, France) (Received February 7th, 1992) 95

Conalbumin-conjugated silica gel, a new chiral stationary phase for high-performance liquid chromatography
by N. Mano and Y. Oda (Ibaraki, Japan), T. Miwa (Gifu, Japan) and N. Asakawa, Y. Yoshida and T. Sato (Ibaraki, Japan) (Received February 12th, 1992) 105

High-performance affinity chromatography of oligonucleotides on nucleic acid analogue immobilized silica gel columns
by E. Yashima, T. Shiiba, T. Sawa, N. Miyauchi and M. Akashi (Kagoshima, Japan) (Received February 5th, 1992) . 111

Practical approach to the miniaturization of chiral separations in liquid chromatography using packed fused-silica capillary columns
by H. Wännman, A. Walhagen and P. Erlandsson (Lund, Sweden) (Received March 16th, 1992) 121

Enantiomeric separation of racemic isoflavanones and related compounds on (+)-poly(triphenylmethyl methacrylate)-coated silica gel
by S. Antus (Budapest, Hungary), R. Bauer (Munich, Germany), Á. Gottsegen (Budapest, Hungary) and H. Lotter, O. Seligmann and H. Wagner (Munich, Germany) (Received February 18th, 192) 133

Separation of oligogalacturonic acids by high-performance gel filtration chromatography on silica gel with diol radical
by J. Naohara (Okayama, Japan) and M. Manabe (Kagawa, Japan) (Received February 6th, 1992) 139

Determination of hydroxylamine in aqueous solutions of pyridinium aldoximes by high-performance liquid chromatography with UV and fluorometric detection
by W. D. Korte (Aberdeen Proving Ground, MD, USA) (Received March 3rd, 1992) 145

(Continued overleaf)

Separation of benz[*f*]isoindole derivatives of amino acid and amino acid amide enantiomers on a β-cyclodextrin-bonded phase
by A. L. L. Duchateau, G. M. P. Heemels, L. W. Maesen and N. K. de Vries (Geleen, Netherlands) (Received February 11th, 1992) . . . 151

High-performance liquid chromatography–thermospray mass spectrometry of gibberellins
by E. B. Hansen, Jr., J. Abian and T. A. Getek (Jefferson, AR, USA), J. S. Choinski, Jr. (Conway, AR, USA) and W. A. Korfmacher (Jefferson, AR, USA) (Received February 7th, 1992) . . . 157

High-performance liquid chromatographic investigations of stillingia oil
by K. Aitzetmüller, Y. Xin, G. Werner and M. Grönheim (Münster, Germany) (Received February 10th, 1992) . . . 165

Gas Chromatography

Comparison of gas chromatographic–mass spectrometric methods for screening of chlorotriazine pesticides in soil
by G. Durand (Barcelona, Spain), P. Gille and D. Fraisse (Vernaison, France) and D. Barceló (Barcelona, Spain) (Received January 3rd, 1992) . . . 175

Solventless determination of caffeine in beverages using solid-phase microextraction with fused-silica fibers
by S. B. Hawthorne and D. J. Miller (Grand Forks, ND, USA) and J. Pawliszyn and C. L. Arthur (Waterloo, Canada) (Received February 11th, 1992) . . . 185

Supercritical Fluid Chromatography

Characterization of a supercritical fluid chromatographic retention process with a large pressure drop by the temporal average density
by X. Zhang and D. E. Martire (Washington, DC, USA) and R. G. Christensen (Gaithersburg, MD, USA) (Received February 27th, 1992) . . . 193

Retention characteristics of high-molecular-weight compounds in capillary supercritical fluid chromatography
by L. D. Giddings, S. V. Olesik and L. A. Pekay (Columbus, OH, USA) and E. Marti (Dublin, OH, USA) (Received March 3rd, 1992) . . . 205

Planar Chromatography

Reversed-phase thin-layer chromatography of diacylglycerols in the presence of silver ions
by V. P. Pchelkin and A. G. Vereshchagin (Moscow, Russia) (Received December 19th, 1991) . . . 213

Fluorescence reactions of inorganic cations heated on a porous glass sheet for thin-layer chromatography
by M. Yoshioka and H. Araki (Osaka, Japan) and M. Seki, T. Miyazaki, T. Utsuki, T. Yaginuma and M. Nakano (Tokyo, Japan) (Received February 21st, 1992) . . . 223

Formation of zinc complexes during chromatography of porphyrins on fluorescent thin-layer plates
by K. Saitoh, C. Kiyohara and N. Suzuki (Miyagi, Japan) (Received March 5th, 1992) . . . 231

Electrophoresis

Theoretical aspects of chiral separation in capillary electrophoresis. I. Initial evaluation of a model
by S. A. C. Wren and R. C. Rowe (Macclesfield, UK) (Received February 12th, 1992) . . . 235

Chromogenic substrate for *endo*-polygalacturonase detection in gels
by O. Markovič, D. Mislovičová, P. Biely and K. Heinrichová (Bratislava, Czechoslovakia) (Received February 26th, 1992) . . . 243

High-performance capillary electrophoretic analysis of chloramphenicol acetyl transferase activity
by J. P. Landers, M. D. Schuchard, M. Subramaniam, T. P. Sismelich and T. C. Spelsberg (Rochester, MN, USA) (Received February 12th, 1992) . . . 247

Separation of sulphonamides and determination of the active ingredients in tablets by micellar electrokinetic capillary chromatography
by Q.-X. Dang (Xian, China) and Z.-P. Sun and D.-K. Ling (Beijing, China) (Received February 6th, 1992) . . . 259

Qualitative analysis of environmental samples for aromatic sulfonic acids by high-performance capillary electrophoresis
by W. C. Brumley (Las Vegas, NV, USA) (Received March 12th, 1992) . . . 267

SHORT COMMUNICATIONS

Column Liquid Chromatography

Multiple ligand applications in high-performance immunoaffinity chromatography
by J. B. Wheatley (Berkeley, CA, USA) (Received April 3rd, 1992) 273

Novel polymeric reagent for synthesizing 9-fluorenylmethoxycarbonyl L-prolinyl derivatives for chiral high-performance liquid chromatography of amino acids
by Z. Zhang, G. Malikin and S. Lam (Bronx, NY, USA) (Received March 30th, 1992) 279

Automated liquid chromatographic determination of ochratoxin A in cereals and animal products using immunoaffinity column clean-up
by M. Sharman, S. MacDonald and J. Gilbert (Norwich, UK) (Received April 3rd, 1992) 285

Gas Chromatography

Automation of a clean-up procedure for determination of trichothecenes in cereals using a charcoal–alumina column
by W. Langseth and P.-E. Clasen (Oslo, Norway) (Received February 24th, 1992) 290

Gas chromatographic–mass spectrometric detection of trace amounts of organic compounds in the intravenous solution Infusio Darrowi
by J. Lesko, T. Jakubik and A. Michalkova (Bratislava, Czechoslovakia) (Received March 3rd, 1992) 294

Electrophoresis

Quantitation of *l*-epinephrine and determination of the *d*-/*l*-epinephrine enantiomer ratio in a pharmaceutical formulation by capillary electrophoresis
by T. E. Peterson and D. Trowbridge (Fort Worth, TX, USA) (Received April 1st, 1992) 298

Author Index 302

Errata 304